arXiv:1303.5375v1 [math.DS] 21 Mar 2013

CONSTRUCTION OF THE DISCRETE HULL FOR THE COMBINATORICS OF A REGULAR PENTAGONAL TILING OF THE PLANE

MARIA RAMIREZ-SOLANO

Abstract
The article "A regular pentagonal tiling of the plane" by Philip L. Bowers and Kenneth Stephenson defines a conformal pentagonal tiling. This is a tiling of the plane with remarkable combinatorial and geometric properties. However, it doesn't have finite local complexity in any usual sense, and therefore we cannot study it with the usual tiling theory. The appeal of the tiling is that all the tiles are conformally regular pentagons. But conformal maps are not allowable under finite local complexity. On the other hand, the tiling can be described completely by its combinatorial data, which rather automatically has finite local complexity. In this paper we give a construction of the discrete hull just from the combinatorial data.

Introduction

An Euclidean tiling of the plane is said to have the finite local complexity property (FLC for short) if for any ball of radius r there is a finite number of patterns of diameter less than r up to a group of motion, usually translation. A tiling has such property if all its tiles are translation copies of a finite number of prototiles and if the tiles match edge to edge. The theory of C^*-algebra and K-theory for aperiodic Euclidean tilings satisfying this property is well-established. Non FLC tilings on the other hand, have their own peculiarities and as a whole are not well-understood. The focus of this article is on non FLC tilings with simple combinatorics, on a conformal pentagonal tiling of the plane described in [18] which was studied by Kenneth Stephenson, Philip L. Bowers, Bill Floyd, and others. The geometry imposed on such tiling has its origins in the theory of Circle Packing. The result is a beautiful tiling, but its geometry prevents us from studying it with the standard tiling tools. In [14], we showed that the FLC property with respect to this geometry, is useless, for it cannot be used to construct a compact topological space. Thus we have to abandon the notion of FLC with respect to the set of conformal isomorphisms that are defined between open subsets of the plane. Instead, we embark in a new quest of investigating the pentagonal tiling from a new perspective, from its combinatorics equipped with a simpler geometry, one where the combinatorics play just as big a role. Such combinatorial tiling we name it K. We will show that K has FLC with respect to the set of isomorphisms that are defined between subcomplexes of K. Such tiling is FLC but is no longer a tiling of the plane. It is a tiling of a surface. It is natural to ask how much of the standard tiling theory can be transferred to this world. This question is the driving force in the development of this work. It turns out that many results of the standard tiling theory can be carried out in a natural way for the pentagonal tiling in the new setting, but they also pose challenges.

An aperiodic FLC Euclidean tiling gives rise to a groupoid and a compact topological space with the action of translation, and so a dynamical system. Such action

---

Supported by the Danish National Research Foundation through the Centre for Symmetry and Deformation (DNRF92), and by the Faculty of Science of the University of Copenhagen.

comes in handy for showing that the classifying space of the groupoid is the topological space. It follows that the K-theory of the topological space is the same as the K-theory of the C^*-algebra of the groupoid. Non FLC tilings, on the other hand, are studied individually, since traditional techniques do not apply even for tilings with simple combinatorics, like the mentioned conformally regular pentagonal tiling of the plane in [18]. For such tiling we can construct a natural groupoid and a compact topological space but it is missing the action. This poses a challenge for proving that the classifying space of the groupoid is the topological space. I raised this question during my talk in the conference "Non standard hierarchical tilings" held in Copenhagen that I co-organized in September 2012.

This work is divided into two sections. The first section provides the basic terminology and results for investigating the combinatorial pentagonal tiling K, which is a CW-complex homeomorphic to the plane and defined via a subdivision rule, and such tiling is shown in Figure 2. Here, we define what we mean with a ball on K, a patch of K, a supertile of K, and so on. In Theorem 1.14 we show that K is FLC with respect to the set of isomorphisms that are defined between subcomplexes of K. Next, we decorate K by decorating its subdivision rule. Subsection 1.4.1 shows that there is only two good possible choices for decorating the subdivision rule, and Proposition 1.26 shows that the chosen decorated subdivision rule induces a unique decoration on K. The purpose of the decorations on K is to destroy local symmetries. Many properties of the decorations of K are summarized in Lemma 1.27. The main result in this section is Proposition 1.29, which shows that we cannot "rotate" a patch of K with respect to any vertex of the patch.

In the second section, we construct the discrete hull Ξ for the tiling, which is a topological space whose elements are basically tilings that look locally the same as the tiling itself. We make distinctions between elements of this space to the level of vertices, hence the use of the word discrete in the name. Equipping it with an ultrametric d' we show it is compact. Moreover, we define a subdivision map on it, which turns out to be continuous, injective, but not surjective. In Theorem 2.14 we show that Ξ is a Cantor space, the main result of this article. A considerable amount of work (and good ideas) goes into finding the right definition of the discrete hull, so that we could construct an étale equivalence relation on it and extend it to a continuous hull - results that will be covered in other articles. The main reason for adding decoration to the tiles is precisely to accomplish this task. Simple as it may seem, the definition of the discrete hull is the fruit of many failed attempts and contradictions found during its construction, some of which are described at the beginning of section two. Elements of the discrete hull are formally isomorphism classes of CW-complexes homeomorphic to the plane, locally isomorphic to K, and with a distinguished vertex. In these attempts, we tried distinguished faces instead of distinguished vertices, or distinguished faces and vertices instead of just distinguished vertices, but we stick to distinguished vertices because in this setting things made sense more quickly, even though another setting could pay off better in the long run.

1. The combinatorial tiling K

This section provides basic terminology and results for investigating the combinatorial pentagonal tiling K introduced in [14] and restated below. Here, we define what we mean with a ball on K, a patch of K, a supertile of K, and so on. In Theorem 1.14 we show that K is FLC with respect to the set of isomorphisms that are defined between subcomplexes of K. Next, we decorate K by decorating its subdivision rule. Subsection 1.4.1 shows that there is only two good possible



Figure 1. Subdivision map.

choices for decorating the subdivision rule, and Proposition 1.26 shows that the chosen decorated subdivision rule induces a unique decoration on K. The purpose of the decorations on K is to destroy local symmetries. The properties of the decorations of K are summarized in Lemma 1.27. The main result in this section is Proposition 1.29, which shows that we cannot "rotate" a patch of K with respect to any vertex of the patch.

We start with some basic definitions. A *combinatorial tiling* is a 2-dimensional CW-complex $(X, \mathcal{E})$, such that X is homeomorphic to the open unit disk $\mathbb{D} := D^2$, where X is a topological space, and $\mathcal{E}$ a partition of X satisfying the CW-complex conditions. Such definition was introduced in [14] and we refer the reader to this article for more details. The *combinatorial tiles* are the closed 2-cells. We also say that a face is a closed 2-cell, an edge a closed 1-cell, and a vertex a 0-cell. A *cell-preserving* map between between CW-complexes is a continuous map between the CW-complexes that maps cells to cells.

Example 1.1. If T is a tiling of the plane, then T has the structure of a 2-dimensional CW-complex, where the 2-cells are the interior of the tiles, the 1-cells are the interior of the edges of the tiles, and the 0-cells are the vertices of the edges of the tiles. Hence, under this identification, $(\mathbb{C}, T)$ is a combinatorial tiling.

Definition 1.2 (subdivision of a combinatorial tiling). Let $(X, \mathcal{E})$ and $(X, \mathcal{E}')$ be two combinatorial tilings with same topological space X. We say that $(X, \mathcal{E}')$ is a subdivision of $(X, \mathcal{E})$ if for each cell $e' \in \mathcal{E}'$, there is a cell $e \in \mathcal{E}$ such that $\overline{e'} \subset \overline{e}$.

Definition 1.3 (subdivision rule). We define the *subdivision rule* ω by Figure 1.

Definition 1.4 (finite CW-complex K_n). Define K_0 as a combinatorial pentagon, which is a space homeomorphic to the closed unit disk with five distinguished points on its boundary. Define $K_n := \omega^n(K_0)$, $n \in \mathbb{N}_0$. Every K_n has a distinguished central pentagon. We define $\iota_n : K_n \to K_{n+1}$ as an embedding which maps the central pentagon of K_n to the central pentagon of K_{n+1}.

Definition 1.5 (the combinatorial tiling K). Define the complex

$$K := \lim_{n\to\infty} K_n,$$

as the direct limit of the sequence of the finite CW-complexes K_n and embeddings ι_n. It has a canonical CW-structure coming from the CW-structure of the complexes K_n, where K_n is obtained from K_{n-1} by attaching finitely many cells. Each cell in the limit K is the image of a cell in K_n for some n.

1.1. *Properties of the ball $B(v, n, K)$*

Any two vertices of K can be joined with finite paths of edges, as K is simply-connected i.e. all its vertices are interior. The length of a path is its number of



FIGURE 2. Balls of different radii but same center.

edges. There exists at least one path between two vertices with smallest number of edges. The distance between two vertices of K is defined as the length of the shortest path between them. Such paths are called *distance-paths*.

DEFINITION 1.6 (Ball $B(v, n, L)$). We define the ball $B(v, n, L)$ as the subcomplex of a combinatorial tiling L whose tiles have the property that all its vertices are within distance n of the vertex $v \in L$.

Since $B(v, n, L)$ is a subcomplex of L, it is closed in L. Examples of balls with same center but different radii are shown in Figure 2. We define the *boundary of the ball* $B(v, n, K)$ to be the edges and vertices satisfying the following condition: if e is an edge of faces f, f' and f is in the ball and f' is not in the ball, then e is on the boundary of the ball.

LEMMA 1.7. *The distance from any boundary vertex of the ball* $B(u, n, K)$ *to the ball's origin* u *can only be* n *or* $n - 1$.

PROOF. By definition of the ball $B(u, n, K)$, where the vertex $u \in K$ and $n \in \mathbb{N}$ are fixed, the distance from any boundary vertex v to u is at most n. Thus we have found an upper bound. Suppose v is a boundary vertex such that the distance to u is $n - 2$. Then we can add in the worse case scenario a pentagon like the one shown in Figure 3(b) to the ball, and so $n - 2$ is not a boundary vertex, a contradiction.



FIGURE 3. (a) Distance to boundary vertex is n. (b) Distance to boundary vertex is n. (c) Distance to boundary vertex is $n-1$.

We will continue investigating the ball $B(u, n, K)$, where the vertex $u \in K$ and $n \in \mathbb{N}$ are fixed. By definition of the ball $B(u, n, K)$, the distance from any boundary vertex v to u is at most n, and thus any $(n+1)$-distanced vertex is automatically outside the ball $B(u, n, K)$. Lemma 1.7 tells us that the distance of the boundary vertices of a ball with respect to the ball's origin is n or $n-1$. A natural question arises. Are all the n and $(n-1)$-distanced vertices from u on the boundary or even in the ball itself? The following lemma shows that there can be n-distanced vertices outside the ball exactly one unit away from the boundary, but all the $(n-1)$-distanced vertices cannot be outside. However, both n and $(n-1)$-distanced vertices can be in the ball's interior. See for example the ball of radius 14 in Figure 2. Observe that we have not discarded the possibility that the ball of radius n has holes. That is, we have not ruled out the possibility that vertices of distance greater than n are surrounded by vertices of distance smaller or equal to n.

One obvious but crucial observation is the following: If v is an n-distanced vertex with respect to vertex u, then v must be neighbor to a $(n-1)$-distanced vertex, else how did we measure such distance?

LEMMA 1.8. *All $(n-1)$-distanced vertices from u are in the ball $B(u, n, K)$. All n-distanced vertices from u are at most one unit away from the boundary of $B(u, n, K)$.*

PROOF. Let v be a $(n-1)$-distanced vertex from u. By definition of vertex-vertex-distance there exists a neighbor to v of distance $n-2$. Then we can attach a pentagon from Figure 3 in worse case scenario, as the edge-diameter of a pentagon is 2 and this pentagon contains vertices of distance $n-1$ and $n-2$. Therefore, the vertex v belongs to the ball $B(u, n, K)$. This proves the first part. Let v be a n-distanced vertex from u. By definition of vertex-vertex distance, it has a neighbor of distance $n-1$, which is in the ball as we have just shown that. Hence v is at most one unit away from the boundary of the ball. See figure 3(c).

We now state some observations in a lemma.

LEMMA 1.9. *Let u be our origin vertex. That is, we will measure distances from u. Let v be a n-distanced vertex. Then*
(a) *All the neighbor vertices of v have distance n, $n-1$ or $n+1$. But there exists at least one neighbor vertex of v of distance $n-1$.*
(b) *The diameter of the ball $B(u, n, K)$ is at most $2n$.*
(c) *A distance-path from u to v goes through at least $n/2$ pentagons and at most $2n$*



Figure 4. The numbers on the vertices represent distance from the center of the ball. Here we see a ball of radius 33 that has holes.

pentagons.
(*d*) *All balls are path-connected and face-connected, but some are not simply connected (i.e. some contain holes). See Figure 4.*

Proof. (*a*) The first part follows by definition of vertex-vertex distance. If v is an n-distanced vertex with respect to vertex u, then v must be neighbor to a $(n-1)$-distanced vertex, else how did we measure such distance?
(*b*) Every vertex in the ball has distance at most n. Hence we can join two vertices in the ball by two distance-paths that go through u and hence their distance is at most $2n$.
(*c*) The diameter of a pentagon is 2. Hence we can advance at most 2 units with every pentagon. Thus we need at least half of them to reach our destination. This proves the first part. To advance one unit we need the edge of a pentagon. So we need at most n pentagons to advance n units. However, when a distance-path goes through a 4-degree vertex, it will go through the vertex of a pentagon but not its edge. Hence we need at most $2n$ pentagons to advance n units.
(*d*) The balls are path connected by definition of the ball. The case scenario in Figure 5 never happens, so the balls are chain connected. Indeed, if v has distance $n-2$ or smaller, then the 4 pentagons are in the ball of radius n; if v has distance $n-1$ or larger, then the shaded pentagon is outside the ball. Figure 4 shows that some balls are not simply connected.

Lemma 1.10. *If the boundary* $\partial B(u,n,K)$ *has length* m*, then the boundary* $\partial B(u,n+1,K)$ *has length at most* $10m$*. In particular a ball of radius* n *has less than* 10^{n-1} *faces.*



FIGURE 5. The integer n inside a pentagon denotes the maximum of the distances of the vertices of the pentagon. A ball of radius n cannot have this pattern on its boundary.



FIGURE 6. For each edge and vertex on a boundary we can attach a pentagon. For example for edge e we attach the blue pentagon. For edge e' we attach the yellow pentagon. For vertex v we attach the green pentagon.



FIGURE 7. Prototiles of K.

PROOF. For every edge and every vertex on the boundary of $B(u, n, K)$ there may exist a pentagon that lies outside the ball. See Figure 6. We consider these pentagons as attached pentagons to $B(u, n, K)$. Then, the boundary of the ball $B(u, n+1, K)$ of radius $n+1$ is contained in the union of the attached pentagons. Let m be the length of the boundary $\partial B(u, n, K)$. Then the number of edges of the attached pentagons is at most $5 \times m + 5 \times m = 10m$. Hence, the length of the boundary $\partial B(u, n+1, K)$ is at most $10m$. A ball of radius 1 has less than 1 faces. So a ball of radius 2 has less than 10 faces. By induction, a ball of radius n has less than 10^{n-1} faces.

By construction, K contains exactly three distinct faces - all completely determined by the degree of their vertices. See Figure 7. The first face t_1 has 0



FIGURE 8. The balls $B(v, 5, K)$, $B(v', 5, K)$ are shown in light color. The balls $B(v, 6, K)$, $B(v', 6, K)$ are shown in red and blue, respectively. The first two are isomorphic, while the last two are not.

four-degree vertices and 5 three-degree vertices. The second face t_2 has 2 four-degree vertices and 3 three-degree vertices. The third face t_3 has 3 four-degree vertices and 2 three-degree vertices.

Since different arrangements of the three faces t_1, t_2, t_3 yield balls of radius n, we expect different balls of radius n. For example, in Figure 8 we show two balls $B(v, 5, K)$, $B(v', 5, K)$ in light red, light green, respectively, and two balls $B(v, 6, K)$, $B(v', 6, K)$, in red and green, respectively. The first two are isomorphic. The latter two are not because they have for example different number of t_3 faces.

A natural question arises. How many distinct balls of radius n(fixed) are there? The following lemma gives a crude estimate.

LEMMA 1.11. *There are finitely many distinct balls of radius n.*

PROOF. By the previous lemma, a ball of radius n has less than 10^n faces. Since there are only three different faces, there are less than 3^{10^n} different balls, a finite number.

We will give nicer properties of these balls later on when we introduce more machinery. For now, this is sufficient to show that K is FLC as is shown next.

1.2. *The combinatorial tiling K is FLC.*

We would like to explain in more detail the FLC property of Euclidean tilings, for then adapt it to our combinatorial tilings. A tiling is said to be FLC if for any fixed radius $r > 0$ and if we look for all patterns in the tiling with diameter less than this r, up to motions in G, there are only finitely many. Often, the group G is just translations. Sometimes it is isometries. For example, the pinwheel tiling of the plane has FLC with respect to $G =$ isometries, but not $G =$ translations.

Let $B_r(x)$ be the closed Euclidean ball of radius r. Let $T(B_r(x))$ be the patch of an Euclidean tiling whose tiles intersect the ball $B_r(x)$. Let $X := \{T(B_r(x)) \mid x \in \mathbb{R}^2\}$. We say that $T(B_r(x)) \sim T(B_r(y))$ if one is a translate of the other, ie. if $T(B_r(x)) = T(B_r(y)) + z$. Then the tiling is FLC if the quotient $X/\sim$ is finite.

DEFINITION 1.12 (patch). A *patch* of a combinatorial tiling is a chain-connected subcomplex of dimension 2 with finitely many cells.

DEFINITION 1.13 (finite local complexity (FLC)). We say that a combinatorial tiling L satisfies the *finite local complexity (FLC)* if for any $r > 0$, there are finitely many patches of edge-diameter less than r up to the set of isomorphisms that are defined between subcomplexes of L.

THEOREM 1.14. *The combinatorial tiling K is FLC.*

PROOF. Let $r > 0$ be given. By Lemma 1.11, there are a finite number of distinct combinatorial balls of radius r in K. Since each of these balls has a finite number of cells, there is a finite number of patches of diameter less than r.

1.3. *Supertiles of K.*

The tiles of K are pentagons. In this subsection, we formalize the definitions of supertiles/superpentagons, flowers, petals, superflower, superpetals, superedges of K. We will see for instance that a superflower is the subdivision of a superpentagon. Results in this section are merely observations in a formalized way. For instance, we prove that the degree of the vertices of K are indeed either of degree 3 or 4. We also show that there are only 3 types of pentagons, who are distinguished by the degree of their vertices; they are depicted in Figure 7. We show that the finite CW-complexes K_n are special in the sense that their "corners" have vertices of degree 3.

If p is a patch of K, then $\omega(p)$ is the CW-complex, where all pentagons of p are subdivided with the subdivision rule shown in Figure 1.

DEFINITION 1.15 (superpentagon). A patch p of K is called a *superpentagon* (of degree m) if $p = \omega^m(q)$, for some $m \in \mathbb{N}_0$ and some pentagon q in K.

The following lemma shows that two superpentagons of same degree are identical except in at most five vertices. Moreover, it shows that the boundary of a superpentagon can be identified with a pentagon.

LEMMA 1.16. *Let p, q be two pentagons of K. Let u_i be the vertices of p, and v_i the vertices of q, where $i = 1, \ldots, 5$. Then $\omega^n(p)\backslash\{u_1, \ldots, u_5\}$ is isomorphic to $\omega^n(q)\backslash\{v_1, \ldots, v_5\}$. That is, their superpentagons of same degree are isomorphic up to the vertices of p and q. If the degree of u_i is the same as the degree of v_i, for all $i = 1, \ldots, 5$, then $\omega^n(q) \cong \omega^n(p)$. In particular the subdivision map leaves unchanged the degree of the vertices of p and q, as it only inserts fresh pentagons. Moreover, the boundary $\partial(\omega^n(p))$ is composed of 5 superedges of same length joined by vertices of degree 3 or 4 (its corners) and the vertices on the superedges are of degree 4. Hence the boundary $\partial(\omega^n(p))$ can be identified with a pentagon.*

PROOF. The two pentagons are identical except that the degree of their vertices might differ, but the degree of these vertices affect only the outside of the pentagons, not the inside, so their inside is identical. Hence the subdivision rule gives the same subdivision for both pentagons. See Figure 9. Since p is a pentagon of K, for every edge of p there is a neighbor pentagon sharing the same edge, and since subdivision is also applied to these neighbor pentagons, the vertices on the boundary are of degree 4 (excluding the vertices of p which are of degree 3 or 4).

We now define a flower for a pentagon.

DEFINITION 1.17 (flower). Let p be a pentagon. Let $(p^0, p^1, p^2, p^3, p^4, p^5) := \omega(p)$, where p^0 is the central pentagon surrounded by five pentagons p^1, p^2, p^3, p^4, p^5, called *petals*. We call $(p^0, p^1, p^2, p^3, p^4, p^5)$ the *flower* of p. See Figure 10.

The following lemma tells the degree of the vertices of the petals of a flower.

LEMMA 1.18. *Let p be a pentagon, and consider its flower $(p^0, p^1, p^2, p^3, p^4, p^5)$. All the vertices of p^0 are of degree 3. The degree of the vertices of p^i, $(i fixed)$ $i = 1, \ldots, 5$ are as follows: two are of degree 3, two are of degree 4 and one is of same degree as a vertex of p. See Figure 10.*



FIGURE 9. The superpentagons $\omega^2(p)$, $\omega^2(q)$ are isomorphic if we ignore the vertices v_i, u_i, $i = 1, \ldots, 5$.



FIGURE 10. The flower of p is $(p^0, p^1, p^2, p^3, p^4, p^5)$. The blue numbers are the degree of the vertices on the flower of p. The variable d_i can take the value 3 or 4.

PROOF. The degree of the vertices of p^i are read directly from Figure 10. We just explain a few details: Since p is a pentagon of K, the vertices of p are of degree 3 or 4. The 4-degree vertices shown in the figure 10 are of this degree because of the following. Since p is a pentagon of K, for every edge of p there is a neighbor pentagon sharing the same edge, and since subdivision is also applied to these neighbor pentagons, the mentioned 4-degree vertices are indeed of degree 4.

The following lemma shows what happens when we apply the subdivision rule to flowers. The result is that the pentagons of the flower turn into identical superpentagons except in at most five vertices.

LEMMA 1.19. *Let p be a pentagon, $n \in \mathbb{N}_0$. Then*

$$\omega(\omega^n(p)) = \omega^n(\omega(p)) = (\omega^n(p^0), \omega^n(p^1), \omega^n(p^2), \omega^n(p^3), \omega^n(p^4), \omega^n(p^5)).$$

Moreover, the superpentagons $\omega^n(p^i)$, $i = 0, \ldots, 5$, are isomorphic with each other if we ignore the vertices on the corners of their boundaries.

PROOF. By definition of ω and of flower of p, we have

$$\omega(\omega(p)) = \omega(p^0, p^1, p^2, p^3, p^4, p^5) = (\omega(p^0), \omega(p^1), \omega(p^2), \omega(p^3), \omega(p^4), \omega(p^5)).$$



Figure 11. Superflower of a superpentagon p.

Hence the first statement. Since p^i, $i = 0, \dots, 5$ are pentagons, we get by Lemma 1.16 that their superpentagons $\omega^n(p^0)$, which are of same degree n, are isomorphic if we exclude the corners of their boundaries $\partial(\omega^n(p^i))$.

Since the boundary of a superpentagon can be identified with a pentagon, it makes sense to generalize the subdivision rule to superpentagons so that instead of replacing pentagons with flowers, we replace superpentagons with superflowers. The previous Lemma 1.19 and Lemma 1.16 allows us to make the following definition.

Definition 1.20 (superflower, superpetal). Let $p = \omega^n(q)$ be a superpentagon, where q is a pentagon. Let $(p^0, p^1, p^2, p^3, p^4, p^5) := \omega(p)$, where $p^0 := \omega^n(q^0)$ is the central superpentagon surrounded by five superpentagons $p^i := \omega^n(q^i)$, $i = 1, \dots, 5$, called *superpetals*. We call $(p^0, p^1, p^2, p^3, p^4, p^5)$ the *superflower* of p. See Figure 11.

The superpentagons composing a superflower are identical except in at most 5 vertices.

The central pentagon K_0 of K is characterized by having all its vertices of degree 3. By definition, $K_n := \omega^n(K_0)$ are superpentagons of K_0 of degree n. The superpentagon $K_{n+1} = \omega(K_n)$ can be seen as the superflower of K_n. Since all the vertices of K_0 are of degree 3, all the vertices on the corners of the boundary ∂K_n are of degree 3, because as we saw earlier, the subdivision map leaves unchanged the degree of such vertices. The superflower of K_n consists of the central superpentagon $K_n^0 = K_n$ and five superpetals located in $K_{n+1} \backslash K_n$. We should mention that K_n is always thought of lying in the center of K.

The following lemma is the generalization of Lemma 1.18 to superflowers, and it concerns the degree of the vertices on the corners of the boundaries of their superpentagons. We have shown earlier that a superpentagon p is identical to each of the superpentagons p^i,$i = 0, \dots, 5$ of the superflower of p, except that the degree of the vertices on the corners of their boundary might not coincide. That is, the subdivision map makes six copies of a superpentagon p while messing around with the corners of ∂p by changing the degree of the vertices on the corners of ∂p.

Lemma 1.21. *Let p be a superpentagon and consider its flower $(p^0, p^1, p^2, p^3, p^4, p^5)$. All the vertices on the corners of ∂p^0 are of degree 3. The degree of the vertices on the corners of ∂p^i, $(ifixed)$ $i = 1, \dots, 5$ are as follows: two are of degree 3, two are of degree 4 and one is of same degree as a vertex on the corner of ∂p. See Figure 10.*
For the special case $p = K_n$, the central superpentagon K_n^0 is not isomorphic to any of the superpetals K_n^i, $i = 1, \dots, 5$. But the superpetals are isomorphic to each other: $K_n^i \cong K_n^j$ for $i, j \in \{1, \dots 5\}$.

Proof. Since p is a superpentagon, there is an integer m and pentagon q such that $p = \omega^m(q)$. By Lemma 1.16 the superpetals $p^i = \omega^m(q^i)$, $i = 0, \dots, 5$ are



FIGURE 12. For a superpentagon p, the superpetals satisfy $\omega(p^i) = \omega(p)^i$.

superpentagons of same degree m. By Lemma 1.16, the subdivision map leaves unchanged the degree of the vertices of the pentagons q^i. Hence the degree of the vertices of the petals q^i are the same as the degree of the vertices on the corners of ∂p^i. This together with Lemma 1.18 proves the first statement.

In particular, since all the vertices of the central pentagon K_0 are of degree 3, all the vertices on the corners of ∂K_n are of degree 3 as well. The central superpentagon $K_n^0 = K_n$ is not isomorphic to any of the superpetals K_n^i, $i = 1, \ldots, 5$ because all the vertices on the corners of ∂K_n^0 are of degree 3, while two of the vertices on the corners of ∂K_n^i are of degree 4. But the superpetals are isomorphic to each other because the degree of the vertices on the corners of ∂K_n^i match those of ∂K_n^j, $j = 1, \ldots, 5$, (i, j fixed).

The following lemma shows what happens when we apply the subdivision map to superpetals of a superflower.

LEMMA 1.22. *Let p be a superpentagon, and consider its superflower. Then $\omega(p^i) = \omega(p)^i$ for $i = 0, \ldots, 5$. See Figure 12.*

PROOF. We have

$$\omega^2(p) = \omega(p^0, p^1, p^2, p^3, p^4, p^5) = (\omega(p^0), \omega(p^1), \omega(p^2), \omega(p^3), \omega(p^4), \omega(p^5)).$$

On the other hand

$$\omega^2(p) = \omega(\omega(p)) = (\omega(p)^0, \omega(p)^1, \omega(p)^2, \omega(p)^3, \omega(p)^4, \omega(p)^5).$$

Hence $\omega(p^i) = \omega(p)^i$. We remark that by Lemma 1.16 given a superpentagon p we can obtain the pentagon q that satisfies $p = w^m(q)$ simply by reading off the degrees of the corners of the superpentagon p. By the same lemma, superpentagons of same degree are identical except at their corners, and by the previous Lemma 1.21 we can distinguish a central superpentagon from a superpetal by the degree of their corners.

We defined the superflower of a superpentagon p using the subdivision rule. The subdivision rule on a superpentagon p simply says to insert a fresh pentagon inside each pentagon of p. The reflection rule on $p = K_n$ described in [18] simply says to



Figure 13. Subdivision with decorations.



Figure 14. Notation of decoration of vertices and edges. Observe the cyclic order in which we write them.

reflect p across each of the superedges of ∂p and identify the two superedges that begin on a corner of ∂p. But this reflection rule is simply Definition 1.20.

The following lemma shows that the petals of the flower of K_n are identical to the superpentagons of the petals K_0^i.

Lemma 1.23. *We have* $K_n^i = \omega^n(K_0^i)$ *for* $i = 0, \ldots, 5$.

Proof. On one side

$$K_{n+1} = \omega(K_n) = (K_n^0, K_n^1, K_n^2, K_n^3, K_n^4, K_n^5).$$

On the other hand,

$$K_{n+1} = \omega^{n+1}(K_0) = \omega^n(\omega(K_0)) = \omega^n(K_0^0, K_0^1, K_0^2, K_0^3, K_0^4, K_0^5) =$$
$$(\omega^n(K_0^0), \omega^n(K_0^1), \omega^n(K_0^2), \omega^n(K_0^3), \omega^n(K_0^4), \omega^n(K_0^5)).$$

Hence $K_n^i = \omega^n(K_0^i)$.

1.4. *Decorating* K.

The combinatorial tiling K defined in 1.5 is not suitable for our purposes. It requires decorations, which are implemented using the decorated subdivision rule shown in Figure 13. We need to address the following questions. Is the decoration well-defined? Which type of decorations the vertices of K can have? Are the local symmetries of K destroyed?

Definition 1.24 (decoration). The *decoration* of each vertex and each edge of K will be written in the notation shown in Figure 14.

Lemma 1.25. *The decorated vertices of the central pentagon* K_0 *are 135, 124, 235, 134, 245 (notice the cyclic order). The decorated edges of the central pentagon* K_0 *are* $(12, 34), (12, 45), (23, 45), (23, 51), (34, 51)$.



FIGURE 15. Decorations of the superedges of K_n.



FIGURE 16. The decoration of K_1 induces a decoration on K_2.



FIGURE 17. The decoration of K_1 is copied into K_1^i.

PROOF. The decoration of each vertex and edge of K_0 is read directly from the figure to the right in Figure 13.

The decoration of the vertices tells us the order of the interior decorations of the pentagons that have in common such vertex, but it does not tell about which pentagon we should start putting the decorations on. Similarly, the decoration of an edge tells about the interior decoration of the pentagons that have in common the edge but it does not tell about which pentagon we should start putting the decorations on. The following lemma shows how the decoration of the boundary ∂K_n looks like.

PROPOSITION 1.26. *The decoration induced by the decorated subdivision map on* K *is well-defined and it is unique. The decoration of the boundary* ∂K_n *is in the pattern shown in Figure 15.*

PROOF. We will decorate K inductively, that is, we will decorate K_{n+1} by knowing the decoration of K_n. For $n = 0$ the decorated subdivision rule tells us how to decorate K_1. For $n = 1$, the decorated subdivision rule tells us how to decorate K_2: the subdivision map identifies vertices of K_1 with vertices of $K_2 = \omega(K_1) = (K_1^0, K_1^1, K_1^2, K_1^3, K_1^4, K_1^5)$, and we decorate such vertices of K_2 according to the decorations of K_1. See Figure 16. We now decorate the superpentagons K_1^i by copying the decorations of K_1, while preserving the already decorated vertices. See Figure 17. Since $K_1^0 = K_1$ are identical, the restriction of the decorated K_2 to K_1 is identical to the decorated K_1. For $n = 3$, the map ω^2 identifies vertices of K_1 with vertices of $K_3 = (K_2^0, K_2^1, K_2^2, K_2^3, K_2^4, K_2^5)$. Decorate such vertices according to the decoration of K_1. We now decorate the superpentagons K_2^i by copying the decorations of K_2, while preserving the already decorated vertices. Since $K_2^0 = K_2$ are identical, the restriction of the decorated K_3 to K_2 is identical to K_2. By induction, the decoration of K is well-defined.

Now to obtain the outside decoration of K_n, we need to look at the outside decoration of K_n^0 lying in the center of K_{n+1}. From Figure 13 we see that the half-decoration 12 of an edge of K_0 gets subdivided into two edges of same half-decoration 12 such that vertex 2 of one edge is joined to vertex 1 of the other edge. We could say that 12 got subdivided into 1212, which is the half-decoration of a superedge of the boundary ∂K_1. By induction on n, the half-decoration 12 gets subdivided into $121212 \cdots 12($ 2^n times), which is the half-decoration of the corresponding superedge of ∂K_n. We conclude the same for the other half-decorations 23, 34, 45, 51 by the same argument. Hence the half-decoration of a superedge of ∂K_n is completely determined by the half-decoration of K_0. Hence, the decoration of ∂K_n is completely determined by the decoration of K_0, which we show in Figure 15. The decorated subdivision induces exactly one decoration on K because there is exactly one exterior decoration on each K_n, $n \in \mathbb{N}_0$.

The following lemma shows the possible decorations of the vertices and edges of K.

Lemma 1.27. *There are five decorations for the 3-degree vertices, for the 4-degree vertices, and for the edges of K. More precisely,*

- *all the decorations of the 3-degree vertices of K are 135, 124, 235, 134, 245 (notice the cyclic order),*
- *all the decorations of the 4-degree vertices of K are* 1234, 1245, 2345, 1235, 1345 *(notice the cyclic order).*
- *all the decorations of the edges of K are* $(12, 34), (12, 45), (23, 45), (23, 51), (34, 51)$.
- *all the decorations of the edges of K are* $(i\ i{+}1, i{+}2\ i{+}3)$ *where* $i = 1, \ldots, 5$, *and* $6 \sim 1$.
- *all the decorations of the edges of K are* $(i\ i{+}1, i{+}3\ i{+}4)$ *where* $i = 1, \ldots, 5$, *and* $6 \sim 1$.

Proof. We have seen in Lemma 1.18 and Figure 10 that the subdivision map inserts a copy of K_0 inside each pentagon. It also subdivides the edges of the original pentagon introducing 4-degree vertices, and it leaves unchanged the degree of the original vertices.

Given the decorated 3-degree vertices of K_n, we get by the decorated subdivision rule that the decorated 3-degree vertices of $K_{n+1} = \omega(K_n)$ are the 3-degree vertices in K_n plus the five decorated 3-degree vertices of K_0(with outside decoration) as the substitution ω inserts a copy of decorated K_0 in every pentagon. Since the decorated 3-degree vertices of K_1 are just the decorated vertices of K_0(with outside decoration), we get by induction on n that all the decorated 3-degree vertices of K_n are the five decorated 3-degree vertices of K_0, which are listed in Lemma 1.25.

The possible decorations of the edges of K_0 are listed in Lemma 1.25, and there are only five of them. Since subdivision of each of these decorated edges simply makes two copies of itself as illustrated in Figure 18 (where the cyclic order of the



FIGURE 18. The numbers i,j=1,...,5 (where 6 is identified with 1) represent the possible decorations of the edge. The decoration of an edge subdivides into two copies of itself and induces a four degree vertex.



FIGURE 19. $(K, v) \neq (K, v')$ as the decoration is not preserved: $\phi(1) = 5 \neq 1$.

decoration of the 4-degree vertices is the reversed cyclic order), and the decorated subdivision rule consists of subdividing the edges of the original pentagons and inserting a copy of the decorated $K_1 \backslash \partial K_1$, there are no more possible decorations. Since the 4-degree vertices of K come from subdividing these edges, and we have only five of them, we have only five possible decorations for the 4-degree vertices of K. See Figure 18.

Let ϕ be a cell-preserving automorphism of K that preserves the decoration. That ϕ preserves the decorations means $\phi(j) = j$, $j = 1, \ldots, 5$. For example, if the interior decorations of K_0 at a vertex $v \in K_0$ is 1 and at another vertex $v' \in K_0$ is $j \neq 1$, then $(K, v) \neq (K, v')$ as the decoration is not preserved: $\phi(1) = j \neq 1$. See Figure 19. Thus the decorations destroy the rotational symmetry of K. We put this result into a lemma.

LEMMA 1.28. *The automorphisms of the decorated K is the identity map.*

PROOF. Let ϕ be a cell-preserving automorphism of K that preserves the decoration. If we forget that ϕ preserves the decoration, then by [18], ϕ is a rotation with respect to the central pentagon or a reflection with respect to the central pentagon and a vertex v. Since the decorated central pentagon has no rotations nor reflections, ϕ must be the identity map.

By the previous lemma, we can tell apart (K, v) from (K, v') whenever $v \neq v'$.



Figure 20. Decoration destroys rotation and reflections.

Proposition 1.29. *Let v be a vertex of K and let P and Q be chain-connected finite subcomplexes containing v. If they are cell-preserving isomorphic and v is mapped to itself then $P = Q$.*

Proof. Let $\phi : P \to Q$ be a cell-preserving isomorphism such that $\phi(v) = v$, and we call v a fixed point. If a decorated tile shares a decorated edge with a neighbor decorated tile, there is no reflection along this edge because all our decorated edges have distinct numbers on both sides. If a decorated tile shares a decorated vertex with a neighbor decorated tile, there is no reflection along this vertex because all our decorated vertices have distinct numbers in their decoration. See Figure 20. Thus since the vertex v is a fixed point of the isomorphism, the decorated faces edges and vertices having in common this vertex are also fixed by ϕ, ie. ϕ is the identity map on the neighbor vertices edges and faces v. Pick one of the fixed tiles of P and call it t. Since the tile t is fixed by ϕ, and there is no reflections along edges nor vertices, t and its neighbors must also be fixed by ϕ, i.e. ϕ is the identity map on the neighbor cells of t. By a finite induction on the neighbors, ϕ is the identity map.

The following theorem is a corollary from the previous proposition.

Theorem 1.30. *The automorphisms of any simply-chain-connected patch P of K are exactly the identity map.*

Proof. Since P is simply-connected, its geometric realization is the closed unit disk. Let ϕ be an automorphism of P. By the Brouwer fixed-point theorem, ϕ has a fixed point x_0. If x_0 lies in the interior of a tile t then the tile is fixed because the cell-preserving automorphisms of any of our decorated tiles are the identity map (we only care about the combinatorial structure i.e. vertices are mapped to vertices and edges to edges and faces to faces). If a decorated tile shares a decorated edge with a neighbor decorated tile, there is no reflection along this edge because all our decorated edges have distinct numbers on both sides. If a decorated tile shares a decorated vertex with a neighbor decorated tile, there is no reflection along this vertex because all our decorated vertices have distinct numbers in their decoration. Thus, since the tile t is fixed by ϕ, and there is no reflections along edges nor vertices, t and its neighbors must also be fixed by ϕ. By induction P is fixed by ϕ, i.e. the only cell-preserving automorphisms of ϕ are the identity. If the fixed point x_0 lies in the interior of an edge, then there is no reflection along the edge, again because the numbers in the decoration of an edge are all distinct. See Figure 20. Thus the decorated edge is fixed by ϕ. If two decorated tiles share this decorated edge, then they are also fixed by ϕ since there is no reflection along a decorated



FIGURE 21. The decorated combinatorial tiling K.

edge. Continuing as before we conclude that ϕ is the identity map in this case as well. If the fixed point x_0 is a vertex, then there is no reflection nor rotation along this vertex again because the numbers in the decoration of a vertex are all distinct. Thus the decorated edges and vertices having in common this vertex are also fixed by ϕ. Continuing as before we conclude that ϕ is the identity map in this case as well.

If a patch is not connected, then it can have more than one automorphism. For example, take two isomorphic connected patches P, Q of K such that they are apart from each other. Let R be the patch $P \cup Q$. Then the automorphism that permutes P and Q is not the identity map.

The decorated combinatorial tiling K is shown in Figure 21. It has eleven prototiles, which are shown in Figure 22. Recall that K without decorations has three prototiles, namely those shown in Figure 7.

1.4.1. *Possible decorated substitutions.* The decorated substitution rule was defined in Figure 13. What if we have defined it instead as ω_i, $i = 1, 2, 3, 4$ shown in Figure



FIGURE 22. The 11 prototiles of decorated K.



FIGURE 23. Possible decorated substitution rules.

23? From this figure, we see that ω_1, ω_2, ω_3 are not suitable because they do not destroy symmetries of some vertices or edges. The substitution ω_4 is the "mirror version" of ω. See Figure 24.

2. The discrete hull Ξ

In the whole text, K will always denote the decorated combinatorial tiling described in the previous section. It consists of decorated two-cells, decorated one-cells, and



FIGURE 24. An alternative decoration of K using the subdivision rule ω_4.

decorated zero-cells along with the data of which is contained in which. A closed two-cell is also called a face, a closed one-cell is called an edge, and a zero-cell is called a vertex. One key feature of K is that it has a central pentagon.

Our goal in this section is to construct a Cantor space of combinatorial tilings that are locally isomorphic to K, which we call the discrete hull Ξ. This space should mimic the discrete hull of an Euclidean tiling. However, combinatorial tilings lack the common frame of reference $\mathbb{C}$, which makes things more complicated. For example, should all tilings isomorphic to each other be considered as different tilings? Of course not, because then a simple relabeling of a tiling would give a different tiling, which looks wrong (i.e if we give all the two-cells of K new names, it should still be considered as K). Should all tilings isomorphic to each other then be considered the same tiling? For example should K rotated by one of its automorphisms be considered the same tiling? The answer is no. The automorphisms on a tiling should be considered different tilings. When looking at pairs (L, u), where u is a vertex of a combinatorial tiling L, we arrived to the following ambiguity: Suppose u and v are two vertices of the central pentagon of K and suppose K was not decorated. Then

(1) $(K, u) = (K, v)$ because every ball at u is the same as at v.
(2) $(K, u) \neq (K, v)$ because we write u and v at the same time on K and so we can distinguish them.

Many other problems arouse, but thankfully, the fix to all these problems is to consider isomorphism classes and the decorated subdivision rule.

As a side note, this complication does not arise in Stephenson's article [18] because they are just studying K and nothing else. We only run into all our problems when we start to think about "things that look locally like K". That is, Stephenson's article does not need to consider isomorphism classes but we do: Suppose

$1, 2, 3, \ldots$ are used as the labels for the vertices of K. In Stephenson's article [18], they only consider automorphisms of K, which translates to using the same labels $1, 2, 3, \ldots$, just in different orders and preserving the combinatorial structure, and therefore there are a finite number of automorphisms (D_5) on K. Our situation is very different. We can construct a new CW-complex L simply by using new labels $a_1, a_2, a_3, \ldots$ and preserving the structure. Then L and K are isomorphic, and in this way we can create infinitely many isomorphisms.

Definition 2.1 (combinatorially isomorphic). Two combinatorial tilings are said to be *combinatorially isomorphic* if they are cell-preserving isomorphic. Thus there is a bijection which preserves all data between the 0-cells, 1-cells and 2-cells.

Definition 2.2 (locally isomorphic). A combinatorial tiling L is *locally isomorphic* to K if for every patch P of L there is a patch Q of K such that P and Q are cell-preserving isomorphic.

Informally, with L is locally isomorphic to K we mean that any finite piece of L appears somewhere in a supertile K_n, $n \in \mathbb{N}_0$. Later on, we will show that the converse also holds, i.e. that any supertile K_n of K lies somewhere in L. We define the set, which we call the discrete hull,

$$\Xi := \{[(L, v)]_{isom} \,|L \text{ is locally isomorphic to } K,\, v \in K^0 \text{ a vertex}\}.$$

That is, for our space Ξ, we consider (K, v) plus all other pairs (L, v) locally isomorphic to it, modulo cell-preserving isomorphism. We denote with $[(L, v)]_{isom}$ the isomorphism class of (L, v). If (L', v') is in $[(L, v)]_{isom}$, then L' is cell-preserving isomorphic to L and the isomorphism identifies v' with v. We say that the vertex u in a pair (L, u) is the origin of L, and we also say that (L, u) is a pointed combinatorial tiling or a combinatorial tiling with origin.

Since any combinatorial tiling is homeomorphic to the plane, and every tiling of the plane is countable, the combinatorial tilings are countable, i.e. has countably many faces.

As a side note we remark that the counterpart notion of aperiodicity of Euclidean tilings is isomorphisms classes. In this sense, the discrete hull is composed of aperiodic tilings.

Lemma 2.3. *The discrete hull Ξ has uncountably many elements.*

Proof. Let $P \subset K$ be a patch. Then $P \subset K_n$ for some $n \in \mathbb{N}$. Since there are infinitely many copies of K_n in K, K has infinitely many copies of P. Since the automorphisms of K are only the identity map, there exist infinitely many copies of P contained in distinct larger patches. Let $P_0 \supset P$, $P_1 \supset P'$ be two such distinct larger patches containing two copies P, P' of P. More precisely, we assume that the interiors of P_0, P_1 contain P, P', respectively, and we can even assume that $P_0 \cap P_1 = \emptyset$. Since the above argument holds for any arbitrary patch P, it holds in particular for our patch P_0. Hence, there exist two distinct patches $P_{00} \supset P_0$, $P_{01} \supset P_0'$ containing two copies P_0, P_0' of P_0. In this way, we obtain the tree of patches,



On such tree, there exist uncountably many paths starting at (P, v) and going down, where v is a fixed vertex of P. Each of these paths yield an element of Ξ simply by superimposing the patches along the path, i.e. by taking the direct limit. Two different paths will yield two different elements of Ξ, as the patches on each row are by construction distinct.

2.1. *The subdivision map ω.*
In this subsection we introduce an injective subdivision map ω on Ξ. Let L be a combinatorial tiling locally isomorphic to K. Define

$$\omega(L) := \bigcup_{t \in L} \omega(t),$$

which is also a combinatorial tiling locally isomorphic to K. If v is a vertex of L, then $\omega(v)$ is a vertex of $\omega(L)$. See Figure 25. We define $\omega(L, v) := (\omega(L), \omega(v))$. If no confusion arises we identify the vertex $\omega(v)$ of $\omega(L)$ with v, and in such case we write $\omega(L, v) := (\omega(L), v)$.

Define the subdivision map $\omega : \Xi \to \Xi$ by

$$\omega([L, v]_{isom}) := [\omega(L), \omega(v)]_{isom}.$$

This map is well-defined, for if $(L, v) \cong (L', v')$ so is $\omega(L, v) \cong \omega(L', v')$.

Theorem 2.4. *The map $\omega : \Xi \to \Xi$ is injective.*

Proof. Suppose that $\omega(L, v) \cong \omega(L', v')$, and let ϕ be the isomorphism. We would like to show that (L, v) is isomorphic to (L', v'). The idea of the proof is that we can recognize (L, v) in $\omega(L, v)$ and (L', v') in $\omega(L', v')$; since there is only one way to do this, the "restriction" of the isomorphism to (L, v) gives the desired result $(L, v) \cong (L', v')$.
Since $\omega(v) \cong \omega(v')$, the map ϕ identifies v with v'. The map ϕ identifies the neighbor vertices of v with the neighbor vertices of v' in a unique way. Indeed, the neighbor vertices of v' are those obtained by (1) start at $\omega(v)$, (2) go along the edges that have $\omega(v)$ as a vertex, (3) ignore the incoming edges from both sides and arrive to a new vertex, which is neighbor to v. See Figure 26. In this way, the map ϕ identifies the neighbor vertices, edges and faces of v with those of v'. Repeating the same argument on the neighbor vertices, ϕ identifies neighbors of neighbors of v with those of v'. By induction, ϕ identifies (L, v) with (L', v'), and thus, they are isomorphic.

Lemma 2.5. *For the combinatorial tiling K we have $\omega(K) \cong K$, but $\omega(K, v) \ncong (K, v)$ for each vertex $v \in K$.*

Proof. By definition of $\omega(K)$, we have $\omega(K) \cong K$. The distance of the central pentagon of K to v is not the same as the distance of the central pentagon of $\omega(K)$ to $\omega(v)$, for any $v \in K$. So $\omega(K, v) \ncong (K, v)$ for any vertex $v \in K$. See Figure 25.



FIGURE 25. The combinatorial tiling (K, v) is shown in red and the combinatorial tiling $(\omega(K), \omega(v))$ is shown in blue. Both have in common the vertex $\omega(v) = v$.



FIGURE 26. Reconstructing (L, v) from $\omega(L, v)$. The neighbor vertex u in L is obtained from two red edges of $\omega(L, v)$ ignoring the incoming green edges of $\omega(L, v)$.



FIGURE 27. The tiling L is defined by $\lim_{n\to\infty} \omega^n(B(v, 2, K))$ with central vertex v, a three degree vertex.



FIGURE 28. The tiling L' is defined by $\lim_{n\to\infty} \omega^n(B(v', 2, K))$ with central vertex v', a four degree vertex.

LEMMA 2.6. *The map* $\omega : \Xi \to \Xi$ *has fixed points.*

PROOF. The tilings (L, v), (L', v') shown in Figure 27 and Figure 28 are fixed points of ω. i.e. $\omega(L, v) \cong (L, v)$, $\omega(L', v') \cong (L', v')$.

2.2. *The metric space* (Ξ, d).
In this subsection, we equip Ξ with an ultrametric d, and we show that (Ξ, d) is a Cantor space. Recall that in Definition 1.6 we defined the ball $B(v, n, L)$ on a combinatorial tiling L.

Definition 2.7 (metric d' on Ξ). Let $d' : \Xi \times \Xi \to [0, \infty)$ be given by

$$d'([L, v]_{isom}, [L', v']_{isom}) := \min(\frac{1}{n}, 1),$$

where $n \in \mathbb{N}$ is the greatest radius where the two balls $B(v, n, L) \simeq B(v', n, L')$ agree assuming the isomorphism maps v to v'.

Notice that $B(v, 0, L) = B(v, 1, L) = \emptyset$, so if the distance between two tilings of Ξ is 1 the two tilings do not agree on any face containing their respective origins, but they might agree on an edge.
We make the following observations. If the distance $1/m = d'((L, v), (L', v')) \leq 1/n$ then $n \leq m$. Informally, this means that we can superimpose (L, v) with (L', v') at their origins v, v', and they will agree on concentric balls of radii $n, \ldots, m$ with center at the origin. If (L, v) and (L', v') agree on a ball of radius n centered at v, v' respectively, then their distance is $d((L, v), (L', v')) \leq 1/n$. If $(L, v), (L', v')$ agree on a ball of radius n with center v, v', respectively, then they agree of course on a ball of radius $n/2$. Hence the distance $d'((L, v), (L', v')) \leq 1/n$ implies $d'((L, v), (L', v')) \leq 2/n$, as expected. The expression $1/(2n) \leq d'((L, v), (L', v')) \leq 1/n$ means that $(L, v), (L', v')$ agree for sure on balls of radii $n, \ldots, 2n$. The expression $1/(2n) < d'((L, v), (L', v')) \leq 1/n$ means that $(L, v), (L', v')$ agree for sure on a ball of radius n, but they do not agree on a ball of radius $2n$.

Recall that $B(v, n, L)$ is a subcomplex of (L, v), and thus the boundary edges and vertices are included in the ball. We also assume that the degree and decoration of the boundary edges and vertices is included in the boundary of the ball.

Lemma 2.8. *Let* $(L, v), (L', v')$ *be two combinatorial tilings locally isomorphic to* K*. If* $B(v, n, L) \cong B(v', n, L')$ *for any integer* n*, then* $(L, v) \cong (L', v')$.

Proof. For short, let $B_n := B(v, n, L)$ and $B'_n := B(v', n, L')$. We have the following inclusions

$$B_1 \subset B_2 \subset \cdots B_n \subset \cdots \subset L,$$

$$B'_1 \subset B'_2 \subset \cdots B'_n \subset \cdots \subset L'$$

and the following isomorphisms $\phi_n : B_n \to B'_n$ satisfying $\phi_n(v) = v'$. Using these maps we need to construct an isomorphism ϕ from (L, v) to (L', v') such that $\phi(v) = v'$. By definition $\phi_n(B_n) = B'_n$ and $\phi_{n+1}(B_n) \cong B'_n$ as combinatorial isomorphisms are isometric but the latter might not be equality. Hence we cannot use all ϕ_n to define ϕ. However, all balls $\phi_n(B_k)$, $n \in \mathbb{N}$ for fixed k are in L' and are isomorphic to B'_k. Since the types of balls of radius k is finite as balls have finitely many cells, a pattern in $\{\phi_n(B_1)\}_{n\in\mathbb{N}}$ must repeat infinitely many times. Thus we can extract a subsequence $\{\phi_{\alpha_1(n)}\}_{n\in\mathbb{N}}$ such that all the balls $\{\phi_{\alpha_1(n)}(B_1)\}_{n\in\mathbb{N}}$ of radius 1 are of the same type. Repeating the same argument, we can extract a subsequence $\{\phi_{\alpha_2\circ\alpha_1(n)}\}$ such that all the balls $\{\phi_{\alpha_2\circ\alpha_1}(B_1)\}_{n\in\mathbb{N}}$ of radius 1 are of the same type and all the balls $\{\phi_{\alpha_2\circ\alpha_1}(B_2)\}_{n\in\mathbb{N}}$ of radius 2 are of the same type. By induction, we can extract a subsequence $\{\phi_{\alpha_k\circ\cdots\circ\alpha_1(n)}\}$ such that it gives balls of same type of radius $1, \ldots, k$. We define ϕ using $\phi_{\alpha_k\circ\cdots\circ\alpha_1(n)}$ choosing k as large as needed.

Theorem 2.9. *The metric* d *on* Ξ *is an ultrametric.*

PROOF. 1) By definition d is positive. 2) We have $d'([L,v]_{isom}, [L,v]_{isom}) = 0$ as (L,v) agrees on itself on any ball of any radius n centered at v.
3) If $d'([L,v]_{isom}, [L',v']_{isom}) = 0$ then $B(v,n,L) \cong B(v',n,L')$ for any integer n, and therefore by the previous lemma $[L,v]_{isom} = [L',v']_{isom}$. 4) By definition, $d'([L,v]_{isom}, [L',v']_{isom}) = 1/n = d'([L',v']_{isom}, [L,v]_{isom})$. 5) It remains to show the ultra triangle inequality: $d'(x,z) \leq \max(d'(x,y), d'(y,z))$. Suppose that $d'(x,y) = 1/n$ and $d'(y,z) = 1/m$. Then x and y agree on a ball of radius n, and y and z agree on a ball of radius m. Hence x and z agree on a ball of radius $\min(x,y)$. Hence $d'(x,z) \leq 1/\min(m,n) = max(1/n, 1/m) = max(d'(x,y), d'(y,z))$. Since $\max(1/n, 1/m) \leq 1/n + 1/m$, an ultrametric is in particular a metric.

LEMMA 2.10. *Let* $\{[L_n, v_n]_{isom}\}_{n\in\mathbb{N}}$ *be a sequence in* Ξ. *Let* $k > 1$ *be such that* $B(v_n, n, L_n) \neq \emptyset$. *If* $B(v_n, n, L_n) \cong B(v_{n+1}, n, L_{n+1})$ *for all* $n \geq k$ *such that* v_n *is mapped to* v_{n+1}*, then there exists a* $[L,v]_{isom} \in \Xi$ *such that* $B(v_{n_i}, n, L_{n_i}) \cong B(v,n,L)$ *for all* $n \in \mathbb{N}$ *with* v_{n_i} *mapped to* v.

PROOF. Define the complex

$$L = \lim_{n\to\infty} B(v_n, n, L_n),$$

as the direct limit of the sequence of balls and isomorphisms. It has a canonical CW-structure coming from the CW-structure of the complexes $B(v_n, n, L_n)$. (The ball $B(v_n, n, L_n)$ is obtained from $B(v_{n-1}, n-1, L_{n-1})$ by attaching finitely many cells.) Each cell in the limit L is the image of a cell in $B(v_n, n, L_n)$ for some n.

LEMMA 2.11. *The (ultra) metric space* (Ξ, d) *is compact.*

PROOF. Let $\{(L_n, v_n)\}_{n\in\mathbb{N}}$ be a sequence in Ξ. We will find a subsequence converging to some $(L,v) \in \Xi$ using a diagonal argument.

For fixed m, there are only finitely many distinct balls of radius m by Lemma 1.11. Since $\{B(v_i, 1, L_i)\}_{i\in\mathbb{N}}$ is an infinite number of balls of radius 1, there is a specific type that repeats infinitely, say $\{B(v_{\phi_1(i)}, 1, L_{\phi_1(i)})\}_{i\in\mathbb{N}}$, where $\phi_1 : \mathbb{N} \to \mathbb{N}$ is a strictly increasing map. Repeating the same argument on the sequence $\{(L_{\phi_1(i)}, v_{\phi_1(i)})\}_{i\in\mathbb{N}}$, we can extract a subsequence

$$\{(L_{\phi_2(\phi_1(i))}, v_{\phi_2(\phi_1(i))})\}_{i\in\mathbb{N}}$$

such that $B(v_{\phi_2(\phi_1(i))}, 2, L_{\phi_2(\phi_1(i))}) \cong B(v_{\phi_2(\phi_1(j))}, 2, L_{\phi_2(\phi_1(j))})$, for all $i,j \in \mathbb{N}$. The map $\phi_2 \circ \phi_1 : \mathbb{N} \to \mathbb{N}$ is a strictly increasing map. By induction we have constructed a subsequence $\{(L_{\phi_n\circ\cdots\circ\phi_1(i)}, v_{\phi_n\circ\cdots\circ\phi_1(i)})\}_{i\in\mathbb{N}}$ containing same type of balls of radius $n, n-1, \ldots, 2, 1$, where $\phi_n \circ \cdots \circ \phi_1 : \mathbb{N} \to \mathbb{N}$ is a strictly increasing map. In particular, $\{(L_{\phi_n\circ\cdots\circ\phi_1(n)}, v_{\phi_n\circ\cdots\circ\phi_1(n)})\}_{n\in\mathbb{N}}$, is a subsequence containing same type of balls of radius $n \in \mathbb{N}$. It is a subsequence because $\phi_{n+1} \circ \cdots \circ \phi_1(n+1) > \phi_n \circ \cdots \circ \phi_1(n)$ as $\phi_n \circ \cdots \circ \phi_1(n+1) > \phi_n \circ \cdots \circ \phi_1(n)$ holds, and so $\phi_{n+1}(\phi_n \circ \cdots \circ \phi_1(n+1)) > \phi_{n+1}(\phi_n \circ \cdots \circ \phi_1(n)) \geq \phi_n \circ \cdots \circ \phi_1(n)$. (The last inequality of type $f(x) \geq x$ is a general property of strictly increasing maps from $\mathbb{N}$ to $\mathbb{N}$). Define $\phi : \mathbb{N} \to \mathbb{N}$ by $\phi(n) := \phi_n \circ \cdots \circ \phi_1(n)$. By Lemma 2.10, there is a $[L,v]_{isom} \in \Xi$ such that $B(v,n,L) \cong B(v_{\phi(n)}, n, L_{\phi(n)})$ for all $n \in \mathbb{N}$ because $(L_{\phi(n)}, v_{\phi(n)})$ has same type of ball of radius n as $(L_{\phi(m)}, v_{\phi(m)})$ for any $m \geq n$, where the centers of the balls are the origins of the pointed combinatorial tilings.

The subsequence $\{[L_{\phi(n)}, v_{\phi(n)}]_{isom}\}_{n\in\mathbb{N}}$ converges to $[L,v]_{isom}$ because given $N \in \mathbb{N}$, for $n \geq N$ we have $d'([L_{\phi(n)}, v_{\phi(n)}]_{isom}, [L,v]_{isom}) \leq \frac{1}{n} \leq \frac{1}{N}$, as both $(L_{\phi(n)}, v_{\phi(n)})$ and (L,v) have in common a ball of radius n centered at $v_{\phi(n)}$ and v respectively.

Since every metric space is Hausdorff, our space (Ξ, d') is Hausdorff. Recall that every metric space is compact if and only if it is complete and totally bounded.

Hence our compact metric space (Ξ, d') is complete and totally bounded. Moreover, (Ξ, d') is a Baire space, as every complete metric space is a Baire space.

Lemma 2.12. *The metric space Ξ is totally disconnected.*

Proof. Let $U \subset \Xi$. Suppose that $x \in U$. Then for any $r \geq 0$ we have $U = (B_r(x) \cap U) \cup (B_r(x)^c \cap U)$. Since the metric d' on Ξ is an ultrametric, the ball $B_r(x)$ is clopen. Hence we have written U as a union of (at most two) open sets in the subspace topology of U. Suppose that U contains at least another point y. Then, we can set $r := d'(x,y)/3$, and so $x \in B_r(x)$ and $y \in B_r(x)^c$. Since U is the union of the nonempty open disjoint sets $(B_r(x) \cap U)$, $(B_r(x)^c \cap U)$ in the subspace topology of U, the only connected sets are the one point sets and the empty set. Hence Ξ is totally disconnected.

Since the space Ξ is compact and totally disconnected, it is a pre-Cantor space.

Lemma 2.13. *For any $[L,v]_{isom} \in \Xi$ and any $n \in \mathbb{N}$ there is a $[L',v']_{isom} \in \Xi$ such that $0 < d'([L,v]_{isom}, [L',v']_{isom}) \leq 1/n$.*

Proof. Let n and (L,v) be given. Since L is locally isomorphic to K, there is a patch in K isomorphic to $B(v,n,L)$. Let $v \in K$ also denote the image of vertex $v \in L$ under the isomorphism map. If $(L,v) \ncong (K,v)$ then $0 < d'([L,v]_{isom}, [K,v]_{isom}) \leq 1/n$. If $(L,v) \cong (K,v)$ then instead of (K,v) use the tiling (L,v) from Figure 27.

Theorem 2.14. *The ultrametric space (Ξ, d') is a Cantor space.*

Proof. The ultrametric space (Ξ, d') is compact by Lemma 2.11. It is totally disconnected by Lemma 2.12. It has no isolated points by the previous Lemma 2.13.

2.3. *More properties of the ball $B(v,n,K)$.*

In this subsection we will give more properties of the balls of K.

Lemma 2.15. *If $\gamma \subset K^1$ is a path of minimal length n, then $\omega(\gamma) \in \omega(K)^1$ is a path of minimal length $2n$.*

Proof. Since each edge is divided into two edges, ω on any path doubles in length. This, together with the fact that the shortest path to reach the endpoints of a subdivided edge is the subdivided edge itself implies that ω on a path of minimal length remains a path of minimal length.



Figure 29. Lengths on a subdivided pentagon.

Lemma 2.16. *For any ball in K, we have*

$$B(\omega(v), 2n-2, \omega(K)) \subset \omega(B(v,n,K)) \subset B(\omega(v), 2n+2, \omega(K)).$$

Proof. We first show that $\omega(B(v,n,K)) \subset B(\omega(v), 2n+2, \omega(K))$. Each vertex of a tile in $B(v,n,K)$ has distance at most n from the center of the ball. If all vertices of a tile are n-distanced, then ω on this tile will give vertices of distance at most $2n+2$. See Figure 29. Thus the vertices of each pentagon in $\omega(B(v,n,K))$ will have distance at most $2n+2$ from $\omega(v)$. Hence $\omega(B(v,n,K)) \subset B(\omega(v), 2n+2, \omega(K))$.

The ball $B(v,n,K)$ contains all vertices of distance at most $n-1$. (Recall that it contains some but not all vertices of distance n). Hence $\omega(B(v,n,K))$ contains all vertices of distance at most $2n-2$ from $\omega(v)$. Hence $\omega(B(v,n,K))$ contains the ball of radius $2n-2$ and center $\omega(v)$.

2.4. *More properties on ω.*

Theorem 2.17. *The map $\omega : \Xi \to \Xi$ is not surjective.*

Proof. Let $\Xi^3 := \{[L,v]_{isom} \in \Xi \mid \text{ degree of } v \text{ is } 3\}$, and let $\Xi^4 := \{[L,v]_{isom} \in \Xi \mid \text{ the degree of } v \text{ is } 4\}$. Suppose that ω is surjective. Then $\Xi = \omega^n(\Xi) = \omega^n(\Xi^3) \cup \omega^n(\Xi^4)$ for any $n \in \mathbb{N}$. Let $u \in K$ be a vertex of degree 3, and $v \in K$ a vertex of degree 4. Then for fixed large integer n, $d'(\omega^n(\Xi^3), [\omega^n(K,u)]_{isom}) \leq 1/n$ and $d'(\omega^n(\Xi^4), [\omega^n(K,v)]_{isom}) \leq 1/n$. Thus, any point in $\Xi = \omega^n(\Xi)$ is close to any of the two specific points $[\omega^n(K,u)]_{isom}$, $[\omega^n(K,v)]_{isom} \in \Xi$. In other words, the two balls $B_{\frac{1}{n}}([\omega^n(K,u)]_{isom}), B_{\frac{1}{n}}([\omega^n(K,v)]_{isom}) \subset \Xi$ are a cover of Ξ, a contradiction.

Theorem 2.18. *The map ω is continuous.*

Proof. By Lemma 2.16, if $d'([L,v]_{isom}, [L',v']_{isom}) = 1/n$ then

$$d'(\omega([L,v]_{isom}), \omega([L',v']_{isom})) \leq \frac{1}{2n-2}.$$

Hence for $n \geq 11$, i.e. for $d'([L,v]_{isom}, [L',v']_{isom}) \leq 1/11$,

$$d'(\omega([L,v]_{isom}), \omega([L',v']_{isom})) \leq 0.55 \cdot d'([L,v]_{isom}, [L',v']_{isom}).$$

Hence d' is continuous.

Acknowledgments.
The results of this paper were obtained during my Ph.D. studies at University of Copenhagen. I would like to express deep gratitude to my supervisor Erik Christensen and Ian F. Putnam whose guidance and support were crucial for the successful completion of this project.

References

1. A. F. Beardon, *A Primer on Riemann Surfaces.* Cambridge University Press, 1984.
2. Jean Bellissard, Riccardo Benedetti, and Jean-Marc Gambaudo, *Spaces of Tilings, Finite Telescopic Approximations and Gap-Labeling.* Commun. Math. Phys. 261, (2006) 1-41.
3. A. Connes, *Non-commutative Geometry.*
Academic Press, San Diego (1994).
4. J. W. Cannon, W. J. Floyd, and W. R. Parry, *Finite subdivision rules.* http://www.math.vt.edu/people/floyd/research/papers/fsr.pdf, (2001).
5. Allen Hatcher, *Algebraic Topology.* Cambridge University Press, 2002.
6. Nicolas Bedaride, Arnaud Hilion, *Geometric realizations of 2-dimensional substitutive tilings.* arXiv:1101.3905 [math.GT], (2011).
7. K. Jänich, S. Levy, *Topology.* Springer-Verlag New York Inc., (1984).
8. J. P. May, *A concise course in Algebraic Topology.* The Univeristy of Chicago Press. Chicago and London, 1999.
9. Shahar Mozes, *Tilings, substitution systems and dynamical systems generated by them.* Journal D'analyse Mathematique, (1989).
10. Ian F. Putnam, *The ordered K-theory of C^*-algebras associated with substitution tilings.* Commun. Math. Phys. 214, (2000) 593-605.

11. Jared E. Anderson and Ian F. Putnam. *Topological Invariants for Substitution Tilings and their Associated C^*-algebras.* Department of Mathematics and Statistics, University of Victoria, Victoria B.C. Canada. (1995) 1-45.
12. Johannes Kellendonk and Ian F. Putnam, *Tilings, C^*-algebras and K-theory.* Directions in mathematical quasicrystals, CRM Monogr. Ser., 13, Amer. Math. Soc., Provicence, RI (2000) 177-206.
13. Jason Peebles, Ian F. Putnam, Ian Zwiers *Minimal Dynamical Systems on the Cantor Set.* Lecture notes, (2011).
14. Maria Ramirez-Solano, *A non FLC regular pentagonal tiling of the plane.* arXiv:1303.2000, 2013.
15. E. Arthur Robinson, Jr., *Symbolic Dynamics and Tilings of* $\mathbb{R}^d$. George Washington University. Washington DC. 1991.
16. Lorenzo Sadun, *Topology of Tiling Spaces.* University Lecture Series Vol. 46, Providence, Rhode Island, 2008.
17. B. Solomyak, *Nonperidicity implies unique composition of self-similar translationally-finite tilings.* Disc. Comp. Geom. 1998.
18. Philip L. Bowers and Kenneth Stephenson, *A "regular" pentagonal tiling of the plane.* Conformal geometry and dynamics. An electronic journal of the American Mathematical Society, (1997) 58-86.

Department of Mathematics, University of Copenhagen, Universitetsparken 5, 2100 København Ø, Denmark.
E-mail: mrs@math.ku.dk